

神経脳生理学部門

1. Onaka T, Luckman SM, Antonijevic I, Palmer JR & Leng G. Involvement of the noradrenergic afferents from the nucleus tractus solitarius to the supraoptic nucleus in oxytocin release after peripheral cholecystokinin octapeptide in the rat. *Neuroscience* **66**, 403-412 (1995).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=7477881>
2. Zhu LL & Onaka T. Involvement of medullary A2 noradrenergic neurons in the activation of oxytocin neurons after conditioned fear stimuli. *Eur J Neurosci* **16**, 2186-2198 (2002).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=12473086>
3. Zhu LL & Onaka T. Facilitative role of prolactin-releasing peptide neurons in oxytocin cell activation after conditioned fear stimuli. *Neuroscience* **118**, 1045-1053 (2003).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=12732249>
4. Onaka T, Ikeda K, Yamashita T & Honda K. Facilitative role of endogenous oxytocin in noradrenaline release in the rat supraoptic nucleus. *Eur J Neurosci* **18**, 3018-3026, (2003).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=14656297>
5. Takayanagi Y, Yoshida M, Bielsky IF, Ross HE, Kawamata M, Onaka T, Yanagisawa T, Kimura T, Matzuk MM, Young LJ & Nishimori K. Pervasive social deficits, but normal parturition, in oxytocin receptor-deficient mice. *Proc Natl Acad Sci USA* **102**, 16096-16101 (2005).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=16249339>
6. Takayanagi Y, Kasahara Y, Onaka T, Takahashi N, Kawada T & Nishimori K. Oxytocin receptor-deficient mice developed late-onset obesity. *NeuroReport* **19**, 951-955 (2008).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=18520999>
7. Takayanagi Y, Matsumoto H, Nakata M, Mera T, Fukusumi S, Hinuma S, Ueta Y, Yada T, Leng G & Onaka T. Endogenous prolactin-releasing peptide regulates food intake in rodents. *J Clinical Invest* **118**, 4014-4024 (2008).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=19033670>

8. Yoshida M, Takayanagi Y, Inoue K, Kimura T, Young L, Onaka T & Nishimori K. Evidence that oxytocin exerts anxiolytic effects via oxytocin receptor expressed in serotonergic neurons in mice. *J Neurosci* **29**, 2259-2271 (2009).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=19228979>
9. Onaka T, Takayanagi Y & Leng G. Metabolic and stress-related roles of prolactin-releasing peptide. *Trends Endocrinol Metab* **21**, 287-293 (2010).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=20122847>
10. Tobin VA, Hashimoto H, Wacker DW, Takayanagi Y, Langnaese K, Caquineau C, Noack J, Landgraf R, Onaka T, Leng G, Meddle SL, Engelmann M & Ludwig M. An intrinsic vasopressin system in the olfactory bulb is involved in social recognition. *Nature* **464**, 413-417 (2010).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=20182426>
11. Onaka T, Takayanagi Y & Yoshida M. Roles of oxytocin neurones in the control of stress, energy metabolism, and social behaviour. *J Neuroendocrinol* **24**, 587-598 (2012).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=22353547>
12. Tobin V, Schwab Y, Lelos N, Onaka T, Pittman QJ & Ludwig M. Expression of exocytosis proteins in rat supraoptic nucleus neurones. *J Neuroendocrinol* **24**, 629-641 (2012).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=21988098>
13. Tachibana M, Kagitani-Shimono K, Mohri I, Yamamoto T, Sanefuji W, Nakamura A, Oishi M, Kimura T, Onaka T, Ozono K & Taniike M. Long-term administration of intranasal oxytocin is a safe and promising therapy for early adolescent boys with autism spectrum disorders. *J Child Adolesc Psychopharmacol* **23**, 123-127 (2013).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=23480321>
14. Yamashita M, Takayanagi Y, Yoshida M, Nishimori K, Kusama M & Onaka T. Involvement of prolactin-releasing peptide in the activation of oxytocin neurones in response to food intake. *J Neuroendocrinol* **25**, 455-465 (2013).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=23363338>
15. Yoshida M, Takayanagi Y & Onaka T. The medial amygdala - medullary PrRP-synthesizing neuron pathway mediates neuroendocrine responses to contextual

conditioned fear in male rodents. *Endocrinology* **155**, 2996-3004 (2014).

<https://www.ncbi.nlm.nih.gov/pubmed/?term=24877622>

16. Nagasawa M, Mitsui S, En S, Ohtani N, Ohta M, Sakuma Y, Onaka T, Mogi K & Kikusui T. Social evolution. Oxytocin-gaze positive loop and the coevolution of human-dog bonds. *Science* **348**, 333-336 (2015).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=25883356>
17. Onaka T, Okabe S, Takayanagi Y & Yoshida M. Noxious or Non-Noxious Inputs to Oxytocin Neurons: Possible Roles in the Control of Behaviors. *Interdisciplinary Information Sciences* **21**, 189-195 (2015).
<https://doi.org/10.4036/iis.2015.B.03>
18. Takayanagi Y, Yoshida M, Takashima A, Takanami K, Yoshida S, Nishimori K, Nishijima I, Sakamoto H, Yamagata T & Onaka T. Activation of supraoptic oxytocin neurons by secretin facilitates social recognition. *Biol Psychiatry* **81**, 243-251 (2017).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=26803341>
19. Nasanbuyan N, Yoshida M, Takayanagi Y, Inutsuka A, Nishimori K, Yamanaka A & Onaka T. Oxytocin-oxytocin receptor systems facilitate social defeat posture in male mice. *Endocrinology* **159**, 763–775 (2018).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=29186377>
20. Onaka T & Takayanagi Y. Role of oxytocin in the control of stress and food intake. *J Neuroendocrinol* **31**, e12700 (2019).
<https://www.ncbi.nlm.nih.gov/pubmed/?term=30786104>